

Metadichol® and CD34 Expression in Umbilical Cord Cells

Raghavan PR*

Nanorx Inc., New York, USA

Abstract

Umbilical cord blood has found use in the clinic for more than 40 years in hematopoietic stem cell transplantation therapies to treat patients with bone marrow diseases or to reconstitute the bone of those cancer patients who had to have theirs wiped out to cure their leukemia or lymphoma. A feature is the presence of CD34 antigen in hematopoietic stem and progenitor cells. These cells can differentiate and are self-renewing, multipotent stem cells that give rise to all blood cells of the immune system and erythrocytes and lymphoid (T cells, B cells, and NK cells) lineages. This study describes increased CD34 gene expression in Umbilical Cord (UC) cells upon treatment with Metadichol which is an inverse agonist of AHR (Aryl Hydrocarbon Receptor). UC cells were subjected to treatment at one picogram, 100 picograms, 1nanogram, 100 nanograms and 1microgram per ml of Metadichol for 72 hrs. Cells treated at 1ng have shown the highest increase in expression of CD34 compared to untreated Control. The cells treated with 1pg, 100 picogram/ml demonstrated the multiplicity of CD34 expression as indicated by peak shift compared to treatment with 1ng, 100 ng, and 1 µg

Keywords: CD 34; Cord blood; Stem cells; Multipotency; Umbilical cord blood; Metadichol, AHR; Aryl Hydrocarbon receptor; VDR; RORC; Inverse agonist; TNF alpha; ICAM1

Introduction

Molecules that modulate adult or embryonic stem cell fate can facilitate the use of stem cell therapies to a multitude of diseases [1]. Well-known characterized adult stem cells are hematopoietic stem cells (HSC) [2]. The expected clinical potential has not materialized due to a lack of defined culture conditions for their expansion [3]. Human umbilical cord blood (CB) is rich in hematopoietic stem cells (HSC) similar to that in bone marrow. CB transplants are increasing annually around the world [4]. This has led to growth of blood banks that store CB and have led to lower risk of graft versus-host disease (GVHD) [5].

The broadest utilization of CB is limited by the low number of HSCs per unit, and most CB units have insufficient stem cells for adults. A lot of work has gone into developing technologies for *ex vivo* expansion of HSCs to enable CB transplants. A low number of hematopoietic stem and progenitor cells in cord blood units limit their widespread use in human transplant protocols.

A feature of hematopoietic stem and progenitor cells is the presence of CD34 antigen cells that are self-renewing, multi-potent stem cells that lead to all blood cells of the immune system and erythrocytes) and lymphoid (T cells, B cells, and NK cells) lineages [6]. The highly specialized cells that arise from HSC are essential in defending the body against infection and disease. Today bone marrow transplantations use stem cells and also make use of their potential for regeneration of damaged tissues. Cord blood cells have a relative long telomere DNA in comparison to their analogs from peripheral blood or bone marrow [7]. Cells from cord blood are capable of hematopoiesis for a longer time. They generate more divisions and produce a more significant number of progeny (daughter cells).

The increase in the expression of CD34 antigen mainly precedes cell differentiation. Transplants of hematopoietic cells have assumed an essential role in the treatment of diseases. They hold much promise for the clinical application involving gene therapy, tolerance induction to facilitate allogeneic or xenogeneic organ transplants. Recent work show the potential of HSC is limitless to generate whole organ systems [8-10].

Boitano and colleagues recently reported a purine derivative SR1 that significantly expands human CB HSCs in culture [11]. The authors screened a chemical library of 100,000 heterocyclic compounds for their ability to grow the numbers of CD34 CB cells in culture. They showed that SR1 supported a fifty-fold increase in CD34 cells. SR1 is an antagonist of the aryl hydrocarbon nuclear receptor protein (AhR), which usually mediates xenobiotic responses but have also been implicated in regulating hematopoietic stem/progenitor cells. Knockdown of the AhR resulted in a sustained proliferation of CD34 cells in culture.

A recent clinical trial of *ex vivo* expanded umbilical cord blood CD34 cells used SR-1 and demonstrated better engraftment and improved early recovery of leukocytes, suggesting that *ex vivo* expansion of HSCs might potentially be achievable [12,13]. Similarly, co-cultures of cord blood CD34 cells with mesenchymal stromal cells led to the expression of CD34 cells by a factor of 30.1, improving the time required for neutrophil engraftment to 15 days, compared to 24 days in recipients who received un-manipulated cord blood CD34 cells [14]. Growing evidence indicates that targeting metabolism and cellular stress for HSC expansion might potentially lead to successful HSC expansion approaches for transplantation therapies in the future [15,16].

Metadichol® [17] is a lipid emulsion of long-chain alcohols, and we have recently shown that it is an inverse agonist of AHR [18] the only one in medical literature. Given Boitano's work that AHR antagonists expand CD34 cells in UBC (umbilical cord cells), our results show that Metadichol which is a safe food-based ingredient can increase expression of CD34 in Umbilical cord blood at picogram levels.

*Corresponding author: Raghavan PR, Nanorx Inc., PO Box 131, Chappaqua, NY 10514, USA, Tel: +1-914-671-0224; E-mail: raghavan@nanorxinc.com

Received January 02, 2018; Accepted January 15, 2018; Published January 29, 2018

Citation: Raghavan PR (2018) Metadichol® and CD34 Expression in Umbilical Cord Cells. J Stem Cell Res Ther 8: 409. doi: [10.4172/2157-7633.1000409](https://doi.org/10.4172/2157-7633.1000409)

Copyright: © 2018 Raghavan PR. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Experimental: Work carried out in Skanda labs, Bangalore, India

Reagents

Antibody: CD34-FITC conjugates: BD Pharmingen

Cell Line: Umbilical Cord (UC) cells were sourced from Hi-Media labs. Germany, they were human Wharton's jelly cord blood cells. They were not frozen and expanded after five passages.

Procedure

Culture 1×10^6 cells in a 6-well plate containing 2 ml of complete media. After 24 h of incubation, cells are treated with 1 pg, 100 pg, 1 ng, 100 ng and 1 µg in serum-free DMEM media and incubated for 72 h. After 72 Hr. of treatment, cells were collected and pelleted cells at 4000

rpm for 5 minutes at room temperature and discard the supernatant. Washed the cell pellet twice with 1X PBS. The cell pellet suspended in 100 µL of Sheath fluid and incubated with CD34-FITC antibody for 20mins in the dark. Post incubation, the cells were once washed with 1X PBS and resuspended in Sheath fluid. The treated and untreated cell populations were determined using FACS Caliber (BD Biosciences, San Jose, CA).

Results

UC cells treated with Metadichol at 1 pg, 100 pg, 1 ng, 100 ng, and 1 µg of for 72 hrs. Cells treated at 1 ng has shown the highest increase in expression of CD34 (Figures 1-4, 82.14%) compared to untreated Control. The cells treated at 1pg, 100pg has demonstrated the multiplicity of CD34 expression as indicated by peak shift (Figures 2




Figure 1: Control plots of UC cells.








Figure 2: UC cells treated with 1 pg of Metadichol.



and 3) compared to treatment with 1 ng, 100 ng, and 1 µg (Figures 5 and 6). The fold increase in CD 34 expression at various concentrations is difficult to calculate as the control cells, the signals for CD 34 look more like background noise. Nonetheless, the CD34 cell expression at 100 pg is seen and is very significant.

Discussion

The results herein complement previously reported efforts to enhance the expression of CD 34 and expansion of HSC [19]. HSCs undergo a massive increase in numbers *in vivo* during the process of hematopoietic reconstitution after stress, such as from infection, lipopolysaccharide challenge, chemotherapy, radiation, or transplantation [20,21], but this still cannot be recapitulated by *ex vivo* expansion approaches. Over the past decade, several novel studies have suggested that HSC expansion *ex vivo* might be feasible has led to a better understanding of specific factors regulating HSC self-renewal *in vivo*, and which have led to novel strategies to increase CD 34 enriched cells in *ex-vivo* [22,23]. Singh et al. [24] showed that aryl hydrocarbon receptor has a role in the regulation of hematopoietic and other stem/progenitor cell and Metadichol an inverse agonist of AHR confirms this finding regulation of Stem Cells.

Also, Vitamin D supplementation also can significantly increase HSPCs *in vitro*. Cortes et al. [25] have pointed out how vitamin D signaling plays a role in this expansion and survival in zebrafish embryos and human umbilical cord blood. They showed that HSPCs respond directly to [1,25] (OH)D₃ stimulation via vitamin D receptor-induced transcriptional activation of the inflammatory cytokine CXCL8. VDR is expressed on human CD34 hematopoietic progenitors [26]. When it comes to bone marrow transplantations, patients can almost never find a donor whose HLA surface proteins match perfectly. However, if the HLA proteins of the donor are too different from those of the recipient, then the cells from the bone marrow transplant attack the recipient's cells and destroy them Graft versus Host Disease" (GVHD). The inability of leukemia and lymphoma and other patients to receive bone marrow transplants is the unavailability of matching bone marrow. Globally, patients are unable to get stem cell transplants needed to combat blood cancers such as leukemia because there is no donor match available. Metadichol being an inverse agonist of both VDR and AHR can exploit multiple pathways in its actions on UBC.

Metadichol® has an advantage over other chemical entities in that it is a food-based ingredient and is entirely devoid of side effects, unlike most synthetic molecules. It has the potential to restore a new functional hematopoietic system in recipients in cases where the patient for example has leukemia. We have shown in two studies an MDS patient (see US patent 9,006,292) and a dengue case study where levels of Platelets and Neutrophils and WBC counts restored to normal levels [27]. Metadichol is uniquely positioned to exploit the power of expressing CD34 and its consequent differentiation and expansion that could be a significant step in offering to all patients who are in need of receiving transplants *ex vivo* or preferably as *in-vivo*.

References

1. Mimeault (2007) Stem cells: a revolution in therapeutics-recent advances in stem cell biology and their therapeutic applications in regenerative medicine and cancer therapies. Clin Pharmacol Ther 82: 252. [\[PubMed\]](#)
2. Kondo M (2003) Biology of hematopoietic stem cells and progenitors: implications for clinical application. Annu Rev Immunol 21:759. [\[PubMed\]](#)
3. Hofmeister CC (2007) Ex vivo expansion of umbilical cord blood stem cells for transplantation: growing knowledge from the hematopoietic niche. Bone Marrow Transplant 39:11. [\[PubMed\]](#)
4. Song Chou (2010) Expansion of Human Cord Blood Hematopoietic Stem Cells for Transplantation, Cell Stem Cell 7: 427-428. [\[PubMed\]](#)
5. Hwang WY (2007) A meta-analysis of unrelated donor umbilical cord blood transplantation versus unrelated donor bone marrow transplantation in adult and pediatric patients. Biol Blood Marrow Transplant. 13: 444-453. [\[PubMed\]](#)
6. Anna H (2015) Characteristics of hematopoietic stem cells of umbilical cord blood Cytotechnology. 67: 387-396. [\[PubMed\]](#)
7. Jackson KA, Mi T, Goodell MA (1999) Hematopoietic potential of stem cells isolated from murine skeletal muscle. Proc Natl Acad Sci USA 96: 14482-14486. [\[PubMed\]](#)
8. Jackson KA (2001) Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells. J Clin Invest 107: 1395-1402. [\[PubMed\]](#)
9. Goodell MA (2001) Stem cells: is there a future in plastics? Curr Opin Cell Biol 13: 662-665.
10. Horwitz EM (1999) Transplantability and therapeutic effects of bone marrow derived mesenchymal cells in children with osteogenesis imperfecta. Nat Med 5: 309-313. [\[PubMed\]](#)
11. Boitano AE (2010) Aryl hydrocarbon receptor antagonists promote the expansion of human hematopoietic stem cells. Science 329: 1345-1348. [\[PubMed\]](#)
12. Fares I (2014) Pyrimidoindole derivatives are agonists of human hematopoietic stem cell self-renewal. Science 345: 1509-1512.
13. Wagner JE (2016) Phase I/II trial of StemRegenin-1 expanded umbilical cord blood hematopoietic stem cells supports testing as a stand-alone graft. Cell Stem Cell 18: 144-155. [\[PubMed\]](#)
14. De Lima M (2012) Cord-blood engraftment with ex vivo mesenchymal-cell coculture. N. Engl. J. Med 367: 2305-2315.
15. Liu X (2015) Maintenance of mouse hematopoietic stem cells ex vivo by reprogramming cellular metabolism. Blood 125: 1562-1565. [\[PubMed\]](#)
16. Vannini N (2016) Specification of hematopoietic stem cell fate via modulation of mitochondrial activity. Nat. Commun 7: 13125. [\[PubMed\]](#)
17. Raghavan PR, US Patents: 8,722,093 (2014); 9,034,383 (2015); 9,006,292 (2015).
18. Raghavan PR (2017) Metadichol® A Novel Inverse Agonist of Aryl Hydrocarbon Receptor (AHR) and NRF2 Inhibitor. J Cancer Sci Ther 9: 661-668.
19. Ito K (2016) Self-renewal of a purified Tie2+ hematopoietic stem cell population relies on mitochondrial clearance. Science 354: 1156-1160. [\[PubMed\]](#)
20. Zhang LX (2013) The effect of CD33 expression on inflammatory response in chronic obstructive pulmonary disease. Immunol Invest 42: 701-710. [\[PubMed\]](#)
21. Zhang LX (2007) Inhibition of CD26 in human cord blood CD34+ cells enhances their engraftment of nonobese diabetic/severe combined immunodeficiency mice. Stem Cells Dev 16: 347. [\[PubMed\]](#)
22. Iscove NN, Nawa K (1997) Hematopoietic stem cells expand during serial transplantation *in vivo* without apparent exhaustion. Curr. Biol 7: 805-808. [\[PubMed\]](#)
23. Takizawa H (2011) Dynamic variation in cycling of hematopoietic stem cells in steady state and inflammation. J. Exp. Med 208: 273-284. [\[PubMed\]](#)
24. Singh KP (2009) The aryl hydrocarbon receptor has a normal function in the regulation of hematopoietic and other stem/progenitor cell populations. Biochemical Pharmacology 77: 577-587. [\[PubMed\]](#)
25. Mauricio Cortes (2016) Developmental Vitamin D Availability Impacts Hematopoietic Stem Cell Production. Cell Reports 17: 458-468.
26. Grande A (2002) Physiological levels of 1α, 25 dihydroxy vitamin D₃ induce the monocytic commitment of CD34+ hematopoietic progenitors. J Leukoc Biol 71: 641-651. [\[PubMed\]](#)
27. Raghavan PR (2016) Inhibition of Dengue and other enveloped viruses by Metadichol®, a novel Nanoemulsion Lipid. J Healing Outcomes 8: 19-25.